

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

STIC-LL

7. NO5/14

From: Canella, Karen
Sent: Wednesday, May 14, 2003 3:05 PM
To: STIC-ILL
Subject: ill order 09/230,955

445790

Art Unit 1642 Location 8E12(mail)

Telephone Number 308-8362

Application Number 09/230,955

- ★ 1. American Journal of Pathology:
1993 Feb, 142(2):403-412
1993, 143(4):1150-1158
1984, 114(3):454-460
1996, 148(3):865-875
1965 Sep, Vol. 44, pp. 280-282
2. Cancer Research, 1993 May 15, 53(10 suppl):2287-2299
3. Cancer epidemiology, biomarkers and Prevention, 1996 Jul, 5(7):549-557
4. Lab Investigation:
1980, 42(1):91-96
1988, 58(2):141-149
5. Gynecol Oncol, 1982, 13(1):58-66
6. International Journal of Gynecological Pathology:
1985, 4(4):300-313
1986, 5(2):151-162
1992, 11(1):24-29
7. Differentiation:
1986, 31(3):191-205
1988, 39(3):185-196
8. Cancer (Phila), 1989, 63(7):1337-1342
9. Cancer Res, 1990, 50(16):5143-5152
10. Virchows Arch B Cell Pathol Incl Mol Pathol, 1987, 54 (2):98-110
11. Acta Histochemica et Cytochemica:
1994, 27(3):251-257
1996, 29(1):51-56
12. Archives of Gynecology and Obstetrics, 1989, 246(4):233-242
13. Clin Lab Med, 1995 Sep, 15(3):727-742
14. Clin Obstet Gynaecol, 1984 Apr, 11 (1):5-23

10555206

Skin Calcium-Binding Protein in Squamous Metaplasia of Human Uterine Cervix

J. H. PAVLOVITCH, A. L. DELEZOIDE,
L. DIDIERJEAN, J. H. SAURAT, and A. PFISTER

From the Laboratoire des Tissus Calcifiés, Hôpital des Enfants-Malades, Paris, France; Laboratoire d'Histologie et Embryologie de la Faculté de Médecine Necker-Enfants-Malades, Paris, France; and Clinique de Dermatologie, Hôpital Cantonal Universitaire, Geneva, Switzerland

The distribution of skin calcium-binding protein in squamous cell metaplasia of human endocervix, in normal human skin, and in ovarian cancer was determined by the immunofluorescence technique. A rabbit antiserum specific to rat SCaBP was characterized by Ouchterlony immunodiffusion and by immunoprecipitation of ^{125}I -labeled SCaBP. The specificity of antibody labeling was demonstrated by using preimmune rabbit serum and SCaBP antiserum competitively absorbed with purified SCaBP. In normal human skin SCaBP was found exclusively in the basal layer cell cytoplasm. This protein was not detected in normal columnar epithelium of endocervix. Epithelial tissues in the zone of transition between the cylindrical epithelium of the endocervical mucosa and the stratified squamous epithelium of the exocervix were obtained from 14 patients with a wide variety of squamous cell metaplasia. In the early stage of metaplasia SCaBP was detected exclu-

sively in the cytoplasm of reserve undifferentiated cells. In the terminal stage of metaplasia the SCaBP was present only in the basal cell layer. SCaBP was found in several layers of dysplastic tissue, and this distribution appeared to be related to the loss of normal maturation of the epithelium. SCaBP was also present in squamous cell carcinoma of endocervix especially in the least differentiated regions of the tumor. No SCaBP was detected in any ovarian cancer cells. These findings are potentially useful as a means of early detection of squamous metaplasia and of distinguishing premalignant anaplastic lesions from those that are benign and non-proliferative. In addition, the presence of SCaBP in tumors derived from metaplastic epithelia and its absence in the ovarian cancer indicate that immunohistochemical search for this protein might be of value in tumors in which an epidermoid origin is a possibility. (Am J Pathol 1984, 114:454-460)

WE HAVE recently demonstrated that a vitamin-D-dependent calcium-binding protein is present in rat skin (SCaBP).¹⁻³ This protein is located in the cytosol of the basal layer cells, which are the proliferative cells of the epidermis.⁴ SCaBP shares many of the properties of other intracellular calcium-binding proteins.⁵ It is a low-molecular-weight protein (12,000 daltons), is acidic in nature, and has an amino acid composition very similar to that of other calcium-binding proteins. It has a high content of phenylalanine, and there are striking homologies in the amino acid sequence between this protein and parvalbumin.³ Like parvalbumin, each molecule binds two calcium ions with high affinity.³ There are, however, differences in the primary structure and calcium-binding properties between SCaBP and other low molecular weight calcium-binding proteins.³ Hence, it is not surprising that antisera raised against SCaBP do not cross-react with intestinal CaBP, calmodulin, parval-

bumin, or the S-100 protein.^{1,2,4,6} The function of SCaBP is not known but it is distributed in the basal layer of all squamous epithelia thus far examined.⁶ These epithelial tissues undergo cell division essentially in the basal layer; and when the basal layer cells rise to suprabasal layer, they lose this capability of division as well as SCaBP content, as judged by immunofluorescence techniques.^{4,6} This association suggested that SCaBP might be related to and be a marker for immature squamous epithelial cells capable of proliferation.

In several pathologic disorders, columnar epitheli-

Supported in part by a grant from the Swiss Foundation for Scientific Research (W-3957082).

Accepted for publication September 26, 1983.

Address reprint request to Dr. J. H. Pavlovitch, Hôpital des Enfants-Malades, Tour Technique 6ème étage, 149, rue de Sèvres, 75743 Paris Cédex 15.

um assumes the characteristics of squamous epithelium, a process termed "epidermization" or "squamous cell metaplasia." This raised the question of whether or not such a metaplastic transformation might be associated with the appearance *de novo* of the SCaBP previously demonstrated in the basal layer of native squamous epithelia.⁶ In the present paper, we report evidence for the presence of SCaBP in the metaplastic lesions of columnar epithelium of the endocervix.

Materials and Methods

Normal human skin and carcinoma of ovary were obtained from surgical specimens. Epithelial tissues in the zone of the endocervical cleft were taken during colposcopic investigation. The biopsies were frozen in liquid nitrogen and mounted on the cryostat chuck. Four-micron sections were cut with a universal microtome for immunohistological incubation. Several adjacent sections were stained with hematoxylin and eosin (H&E). The presence of stratified or epidermoid epithelium in the tunnel or clefts of endocervix was considered as a state of epidermization or squamous cell metaplasia. Morphologic criteria were used to distinguish between various degrees of atypism in the case of dysplasia: the detection of "reserve" or undifferentiated basal type cells, uniform in type rather than stratified, was interpreted as an initial stage of metaplastic lesion or "reserve" cell hyperplasia. The presence of multilayered patterns with a morphologic characteristic of stratified squamous epi-

thelia was considered the terminal stage of metaplasia. The multilayered pattern with the presence of variably atypical basal type cells in the upper layers was considered dysplasia. Total or partial cell replacement was taken as the criterion of the gravity of lesion. In cases of mild atypism the atypical cells extended about one-half of the thickness of the epithelial layer. In those of marked atypism these cells penetrated through 75-90% of the epithelial thickness. In 2 patients, the histologic pattern corresponded to an epidermoid variety of cervical cancer (squamous cell carcinoma). In 4 patients selected for this study, the tissue sections of endocervix containing normal columnar epithelium served as controls.

Antiserum

A rabbit antiserum containing antibodies to rat SCaBP was obtained by methods previously described.¹ Rat skin samples were frozen in liquid nitrogen, pulverized in a Spex Mill at -80 C (Spec Inc., Metuchen, NJ), extracted with Tris buffer (1.37×10^{-2} M Tris, 0.119 NaCl, 4.74×10^{-3} M KCl, pH 7.4) and chromatographed on a Sephadex G 75 column. The fractions were analyzed for protein⁶ and for calcium-binding activity by the Chelex 100 competitive binding assay.⁸ Further purification was obtained using DEAR Sephadex A 25 ion exchange and Sephacril G 200 chromatography. After purification, SCaBP produced one band on dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) in the region of 12,000 daltons.² An antiserum to the purified pro-

Table 1—Skin Calcium-Binding Protein Immunofluorescence in Native and Metaplastic Squamous Epithelia

Organs	Histology	Number of specimens examined	Number of SCaBP-positive specimens	Localization of SCaBP staining
Skin	Normal epidermis	50	50	Basal cell layer
	Normal columnar epithelium	4	0	
	Undifferentiated "reserve" cell hyperplasia	2	2	Undifferentiated cell layer
	Terminal stage of metaplasia	4	4	Basal cell layer
	Dysplasia with mild atypism	5	5	Basal and supra-basal cell layer, parallel to penetration of immature cells well
	Dysplasia with marked atypism	1	1	In all cellular layers
	Squamous cell carcinoma	4	4	In tumor cells mainly in the periphery of the tumoral buds
Ovary	Undifferentiated small-cell carcinoma	1	0	




Figure 1 — Human skin. a — Epidermal basal cell layer of epidermis stained with SCaBP antiserum (diluted 1:40). ($\times 625$) (With a photographic reduction of 38%) b — Control section treated with preimmune rabbit serum (serum diluted 1:40) ($\times 250$) (With a photographic reduction of 31%)

tein was raised by repeated injection of white New Zealand rabbits with 50–100 μ g of SCA₂BP. The antiserum had been previously tested by the Ouchterlony double immunodiffusion technique. The antibody precipitation technique was also used to determine the specificity of the antiserum, as previously described.⁶ Cross-reactivity between rat and human SCA₂BP was tested by radioimmunoassay measurement of SCA₂BP in man by the use of rat antibody (unpublished) and by the immunoblotting technique, in which the rat antibody revealed the identical single band for SCA₂BP from rat and human skin (Siegenthaler, et al, submitted for publication).

For the control absorption, the antiserum against SCaBP was incubated with excess purified SCaBP or the total soluble protein extract of skin for 1 hour at 37 C and 12 hours at 4 C. After centrifugation at 2700g for 1 hour the supernatant was taken and used as a reagent for the indirect immunofluorescence procedure.





Figure 2—Human uterine cervix. Hyperplasia of undifferentiated reserve cells. **a**—H&E staining. The hyperplastic "reserve" cells of basal type (*R*) forcing columnar epithelium upward toward the luminal area of cervical canal. ($\times 300$) (With a photographic reduction of 37%) **b**—Section of the same region. Immunofluorescence staining due to anti-ScABP serum. ($\times 625$) Specific fluorescence is seen exclusively in the hyperplastic reserve cell population (*arrow*). No specific fluorescence is observed in the columnar epithelium of endocervix. (With a photographic reduction of 44%) **c**—Control section treated with preimmune rabbit serum. ($\times 625$) No fluorescence in the proliferative reserve cells is observed. (With a photographic reduction of 18%)



Figure 3
details of
treated
of metal
preimm

Ouc
the tes
1) bra
Labou
bourg
Dr. D
Macro
rat in
March

Indire
The
mined
insuff





Figure 3—Human uterine cervix. Terminal stage of metaplastic transformation. **a**—H&E-stained section ($\times 300$) showing the morphologic details of squamous epithelium. Basal layer cell; DP, papilla. (With a photographic reduction of 23%) **b**—Section of the same region treated with SCaBP antiserum. ($\times 625$). Immunofluorescent staining is seen in the basal layer cells (b) of newly formed squamous epithelia of metaplasia; fluorescence in the basal layer cells of papilla (P). (With a photographic reduction of 20%) **c**—Control section treated with preimmune rabbit serum ($\times 625$). Notice the absence of fluorescence in the basal layer. (With a photographic reduction of 20%)

Ouchterlony double immunodiffusion was used for the testing of cross-reactivity. The antigens used were 1) brain S-100 calcium-binding protein (gift of Dr. Labourdette, Laboratoire de Neurochimie, Strasbourg, France) and 2) ram testis calmodulin (gift of Dr. Demaille, Centre de Recherche de Biochimie Macromoléculaire, Montpellier, France). The purified rat intestinal SCaBP was prepared as described by March et al.⁹

Indirect Immunofluorescence

The presence of SCaBP in endocervix was determined only by indirect immunofluorescence,⁴ because insufficient biopsy tissue was obtained for the immu-

noprecipitation technique or radioimmunoassay. For the indirect immunofluorescent test, the tissue sections were incubated with the SCaBP antiserum (diluted 1:10 to 1:640 in phosphate-buffered saline (PBS) for 30 minutes at room temperature). After washing for 30 minutes in PBS, tissues were incubated for 30 minutes with commercially prepared fluorescein isothiocyanate (FITC)-labeled sheep anti-rabbit IgG (H&L) antiserum (Hoescht-Boehringer, Paris, France): specific antibody concentration, 1 mg/ml; FITC protein ratio, 2.2; working dilution, 1:20 in PBS to obtain 50 μ l/ml of specific antibodies. The slides were then washed for 30 minutes in PBS and mounted with glycerol. They were examined with a Leitz Orthoplan microscope equipped for incident illumination. Con-

ated re-
cells of
luminal
of 37%)
due to
lusively
fluores-
x. (With
ted with
ferative
if 18%)




Figure 4—Human uterine cervix. Dysplasia with marked atypia (full thickness lesion). **a**—H&E-stained section showing uniformly small immature cells with a high nuclear/cytoplasmic ratio ($\times 300$). (With a photographic reduction of 5%) **b**—Section of a similar region treated with SCAbP antiserum ($\times 250$). Notice the immunospecific fluorescent staining in all cellular layers (arrow). (With a photographic reduction of 12%) **c**—Control section treated with preimmune rabbit serum ($\times 625$). No fluorescence is observed. (With a photographic reduction of 17%)

trol sections were stained with preimmune serum from the corresponding rabbit or with antiserum against SCAbP absorbed with the antigen as described above.

In order to avoid the artificial migration of water-soluble antigen, the freeze-drying method of fixation was used for control as previously described.⁴

Results

SDS-PAGE of immunoprecipitated ^{125}I counts co-migrate with the ^{125}I -SCaBP marker, showing a single band of radioactivity corresponding to a molecular weight of 12,000. In Ouchterlony double immuno-


diffusion the antiserum against SCAbP recognized the antigen from protein extract of homogenized skin. No precipitation lines were observed when SCAbP antiserum was incubated with protein extract of muscle, intestine, liver, bone, or kidney. SCAbP antiserum did not cross-react with ram testis calmodulin, brain S-100, or intestinal CaBP (not shown).^{4,6}

The location of SCAbP of native and metaplastic squamous epithelium is shown in Table 1. In the normal human skin anti-SCaBP serum gave bright staining of the epidermal basal cell layer (Figure 1). In agreement with our earlier observation in the rat,⁶ the pattern of basal cell immunoreactivity in human skin was consistent with a cytoplasmic staining. The dis-

March 1984

erine cervix.
ed atypism
on). a—
n showing
ature cells
ytoplasmic
a photo-
5%) b—
ilar region
antiserum.
mmunospe-
ining in all
n). (With a
ion of 17%)
reated with
um (x 625).
observed.
reduction.

Figure 5—Human uterine cervix. Squamous cell carcinoma. a—Section treated with SCaBP antiserum. (x 625) Notice the immunofluorescence in tumor cells mainly in the periphery of the tumor buds. s, stroma; t, tumor bud. b—Control section treated with preimmune rabbit serum. (x 250) No fluorescence is observed.



tribution of the antigen in the tissue prepared by freeze-drying was similar to that obtained using the nondried freezing technique. The same distribution, but with lower intensity of specific immunofluorescence, was seen in methanol-fixed sections. No specific SCaBP antibody immunofluorescent staining occurred with SCaBP antiserum that had been competitively absorbed with purified protein or with total protein extract from rat skin.

The normal endocervical tissue did not contain SCaBP. In all cases, zones of squamous metaplasia contained SCaBP antigen. The cytologic distribution of SCaBP, as detected immunohistochemically, was variable. In the case of the initial state of squamous metaplasia, the SCaBP was detected only in the cytoplasm of the hyperplastic undifferentiated cells (Figure 2). In the terminal stage of metaplasia, the specific fluorescence was seen exclusively in the cytoplasm of the basal layer of the metaplastic region (Figure 3).

On the surface of all dysplastic lesions, the SCaBP

immunoreactivity was detected in the small uniform, immature cells. The distribution of SCaBP was parallel to the penetration of this immature type of cell well into the upper layer of epithelium. In the samples considered as having dysplasia with mild atypism, the SCaBP was demonstrated only in the lower layer of immature cells. No staining was observed in the upper layers. In the state of dysplasia with marked atypism ("full thickness" alteration), SCaBP was demonstrated in all cellular layers (Figure 4).

In squamous cell carcinoma SCaBP immunofluorescence was present exclusively in the tumor cell population (Figure 5). No immunoreactivity was observed in the stroma. The percentage of stained cells varied in the different areas of the lesion. The majority of the SCaBP-positive cells was present in the periphery of the tumoral buds in the zones that may be considered histologically as least differentiated. However, several SCaBP-positive cell agglomerations were present in the areas that were histologically more

ized the
ed skin.

SCaBP
tract of
3P anti-
iodulin,
4,6

aplastic
the nor-
stain-
e 1). In
rat,⁶ the
ian skin
the dis-

differentiated. No SCaBP immunoreactivity was present in any region of undifferentiated ovarian cancer.

Discussion

SCaBP, a calcium-binding protein originally isolated from rat skin is distributed exclusively in the undifferentiated basal cells of squamous epithelia.⁶ The present study made in human tissue confirms the earlier observations in rats that SCaBP is present in normal squamous but not in normal columnar epithelia. Moreover, in the case where columnar epithelium assumes the characteristics of squamous epithelium, the *de novo* appearance of SCaBP takes place. This protein may be, therefore, regarded as a common marker of squamous transformation. This finding may have immediate relevance to the detection of the early lesions of squamous metaplasia, especially those in which the morphologic picture is confusing.

The cytologic distribution of SCaBP in the variety of metaplastic lesions of endocervix suggests that its location depends on the degree of squamous epithelial cell maturity. In the terminal state of metaplasia the SCaBP is present only in the basal layer cells, but the multilayered pattern is found in the state of dysplasia, related to loss of normal maturation of the epithelium. This finding may be useful as a means for differentiating premalignant anaplastic lesions from those that are benign and nonproliferative. In addition, the presence of SCaBP in tumors derived from metaplastic epithelia and its absence in ovarian cancer indicate that the immunohistochemical search for this protein might be of value in tumors in which an epidermoid origin is in question.

It is possible that the presence of SCaBP in basal cells of either native or metaplastic stratified epithelia reflects a specific aspect of its function in the maintenance of the undifferentiated proliferative cells in the basal layer of squamous epithelium. It is interesting to note that epidermal and metaplastic cervical epithelia have an unusual response to changes in extracellular calcium concentrations.^{10,11} In contrast to all other cell types examined, which cease to replicate when extracellular calcium levels are decreased below 0.1 M,¹² these cells proliferate only at low calcium concentrations. Higher calcium levels in the culture medium inhibit their mitotic activity and induce stratification and squamous differentiation. It is tempting to speculate that there is a causal relationship between the appearance of SCaBP and the cellular response to calcium concentrations. If this were so, then the metaplastic transformation of cervical epithelium may involve altered cellular calcium metabolism.

Finally, several lines of evidence indicate that many

neoplastic cells *in vitro* do not respond by modification in mitotic rate to changes in extracellular calcium concentrations, in contrast to normal cells.^{12,13} It has also been shown that the regulation of growth and differentiation by calcium is altered in malignant epidermal cells and preneoplastic cells exposed to carcinogenesis.¹⁴ If the SCaBP expression of SCaBP is altered in aberrant differentiation states of tumor cells, as suggested by our results, this protein could be linked to decreased dependence of neoplastic cells on extracellular calcium for growth regulation.

References

1. Pavlovitch JH, Laouari D, Didierjean L, Saurat JH, Balsan S: Vitamin D dependent calcium binding protein in rat epiderm, *Calcium Binding Proteins: Structure and Function*. Amsterdam, Oxford, Elsevier, 1982, pp 417-421
2. Laouari D, Pavlovitch JH, Deceneux G, Balsan S: Vitamin-D dependent calcium-binding protein in rat skin. *FEBS Lett* 1980, 111:285-89
3. Rinaldi ML, Halech J, Pavlovitch JH, Ferraz C, Derancourt J, Demaille JG: Isolation and characterization of a rat skin parvalbumin-like calcium-binding protein. *Biochemistry* 1982, 21:4805-4810
4. Saurat JH, Didierjean L, Pavlovitch JH, Laouari D, Balsan S: Skin calcium-binding protein is localized in the cytoplasm of the epidermis. *J Invest Dermatol* 1982, 76:221-223
5. Kretsinger RH: Calcium-binding protein. *Ann Rev Biochem* 1976; 45:239-246
6. Pavlovitch JH, Didierjean L, Rizk M, Balsan S, Saurat JH: Skin calcium binding protein: Distribution in other tissues. *Am J Physiol* 1983, 244:C50-C57
7. Lowry CH, Rosebrough NJ, Farr AL, Randal RJ: Protein measurement with the folin phenol reagent. *J Biol Chem* 1951, 193:265-275
8. Hermsdorf CL, Brenner F: Vitamin D-dependent calcium binding protein from rat kidney. *Biochem Biophys Acta* 1975, 379:553-561
9. Marche P, Pradelles P, Gros C, Thomasset M: Radioimmunoassay for a vitamin D-dependent calcium-binding protein in rat duodenal mucosa. *Biochem Biophys Res Commun* 1977, 76:1020-1026
10. Henning H, Michael DS, Cheng K, Steinert K, Holbrook K, Yuspa SH: Calcium regulation of growth and differentiation of mouse epidermal cells in culture. *Cell* 1980, 19:245-254
11. Auesperg N, Swierenga SHH, Feng S: Calcium-dependent morphogenesis in a cell line derived from human uterine cervical carcinoma. *Cell Biol Invest Rep* 1982, 6:495-501
12. Swierenga SHH, Whitfield JF, Boynton AL, MacManus JP, Rixon RH, Sikorska M, Tsang BK, Walker PR: Regulation of proliferation of normal and neoplastic rat liver cells by calcium and cyclic AMP. *Ann NY Acad Sci* 1980; 349:294-311
13. Durham CH, Walton JM: Calcium ion and the control of proliferation in normal and cancer cells. *Bioscience Rep* 1982, 2:15-30
14. Yuspa SH, Morgan DL: Mouse skin cells resistant to terminal differentiation associated with initiation of carcinogenesis. *Nature (Lond)* 1981, 293:72-74

Acknowledgments

We thank Dr R. Ouzilu from Clamart Hospital for providing colposcopic samples.

Ultra
Pigs

Large d
ionopho
skeletal
For a stu
in affect
given 48
euthanat
and myo
ropsy, w
muscle
shoulder
necrosis
fibers. Is
contract
amorph
the pers
tubes);

MONE
in 1967
cinnam
ized as
of its a
that ca
monent
subsequ
coccidi
was effe
its acti
of prog
occurre
monem
is used
Seven
in cat
pound
sheep,
turkey
found